

Block numbers, 321-avoidance and Schur-positivity

Eli Bagno (Jerusalem College of Technology)
joint work with Ron M. Adin and Yuval Roichman

Permutation Patterns 2017

Short description of results

We present here three results concerning the **block number** statistic on **321-avoiding permutations**:

- **Equi-distribution** of block number and the complement of last descent over certain sets of 321-avoiding permutations.
- The set of 321-avoiding permutations with a given block number is **symmetric and Schur-positive**.
- An explicit formula for the corresponding **character**.

Outline

- 1 Introduction
- 2 Equi-distribution
- 3 Symmetry and Schur-positivity
- 4 Proof idea
- 5 Open problems

Introduction

Describing pattern-avoiding classes

Let $S_n(\Pi)$ be the set of permutations in S_n avoiding a given set of patterns Π . There are several ways to provide information about this set.

- 1 Compute the cardinality $|S_n(\Pi)|$ (Simion, Wilf, ...).
- 2 Compute the generating function for a statistic *stat*:

$$\sum_{\pi \in S_n(\Pi)} q^{\text{stat}(\pi)}$$

(Sagan, Pak, Elizalde, ...).

- 3 Compute the quasi-symmetric function

$$\sum_{\pi \in S_n(\Pi)} F_{\pi}(x_1, x_2, \dots)$$

(Sagan, Woo, ...).

Describing pattern-avoiding classes

Let $S_n(\Pi)$ be the set of permutations in S_n avoiding a given set of patterns Π . There are several ways to provide information about this set.

- 1 Compute the cardinality $|S_n(\Pi)|$ (Simion, Wilf, ...).
- 2 Compute the generating function for a statistic *stat*:

$$\sum_{\pi \in S_n(\Pi)} q^{\text{stat}(\pi)}$$

(Sagan, Pak, Elizalde, ...).

- 3 Compute the quasi-symmetric function

$$\sum_{\pi \in S_n(\Pi)} F_{\pi}(x_1, x_2, \dots)$$

(Sagan, Woo, ...).

Describing pattern-avoiding classes

Let $S_n(\Pi)$ be the set of permutations in S_n avoiding a given set of patterns Π . There are several ways to provide information about this set.

- 1 Compute the cardinality $|S_n(\Pi)|$ (Simion, Wilf, ...).
- 2 Compute the generating function for a statistic *stat*:

$$\sum_{\pi \in S_n(\Pi)} q^{\text{stat}(\pi)}$$

(Sagan, Pak, Elizalde, ...).

- 3 Compute the quasi-symmetric function

$$\sum_{\pi \in S_n(\Pi)} F_{\pi}(x_1, x_2, \dots)$$

(Sagan, Woo, ...).

Quasi-symmetric functions

Quasi-symmetric functions were defined by Gessel ('84).

Every subset $J \subseteq [n - 1]$ has an associated **fundamental quasi-symmetric function** $F_J(\mathbf{x})$ (to be defined later).

For a set of permutations $A \subseteq \mathcal{S}_n$ define

$$Q(A) = \sum_{\pi \in A} F_{\text{Des}(\pi)}.$$

Symmetry and Schur-positivity

Question (Gessel and Reutenauer, '93)

For which $A \subseteq \mathcal{S}_n$ is $Q(A)$ a symmetric function?

A symmetric function is **Schur-positive** if all the coefficients in its expression as a linear combination of Schur functions are non-negative.

Call $A \subseteq \mathcal{S}_n$ Schur-positive if $Q(A)$ is.

For example,

Theorem (Gessel and Reutenauer, '93)

Conjugacy classes are symmetric and Schur-positive.

Symmetry and Schur-positivity

Question (Gessel and Reutenauer, '93)

For which $A \subseteq \mathcal{S}_n$ is $Q(A)$ a symmetric function?

A symmetric function is **Schur-positive** if all the coefficients in its expression as a linear combination of Schur functions are non-negative.

Call $A \subseteq \mathcal{S}_n$ Schur-positive if $Q(A)$ is.

For example,

Theorem (Gessel and Reutenauer, '93)

Conjugacy classes are symmetric and Schur-positive.

Symmetry and Schur-positivity

Question (Gessel and Reutenauer, '93)

For which $A \subseteq \mathcal{S}_n$ is $Q(A)$ a symmetric function?

A symmetric function is **Schur-positive** if all the coefficients in its expression as a linear combination of Schur functions are non-negative.

Call $A \subseteq \mathcal{S}_n$ Schur-positive if $Q(A)$ is.

For example,

Theorem (Gessel and Reutenauer, '93)

Conjugacy classes are symmetric and Schur-positive.

Symmetry and Schur-positivity

Classical examples of (symmetric and) Schur-positive sets of permutations include:

- Conjugacy classes
- Inverse descent classes
- Knuth classes
- Permutations with a fixed inversion number
- Arc permutations

Problem (Sagan and Woo, '14)

Find sets of patterns Π and parameters $stat$ such that $Q(\{\sigma \in \mathcal{S}_n(\Pi) \mid stat(\sigma) = k\})$ is symmetric and Schur-positive.

Symmetry and Schur-positivity

Classical examples of (symmetric and) Schur-positive sets of permutations include:

- Conjugacy classes
- Inverse descent classes
- Knuth classes
- Permutations with a fixed inversion number
- Arc permutations

Problem (Sagan and Woo, '14)

Find sets of patterns Π and parameters $stat$ such that $Q(\{\sigma \in \mathcal{S}_n(\Pi) \mid stat(\sigma) = k\})$ is symmetric and Schur-positive.

Equi-distribution

Direct sum of permutations

Definition

Let $\pi \in \mathcal{S}_m$ and $\sigma \in \mathcal{S}_n$. The **direct sum** of π and σ is the permutation $\pi \oplus \sigma \in \mathcal{S}_{m+n}$ defined by

$$(\pi \oplus \sigma)_i = \begin{cases} \pi(i), & \text{if } i \leq n; \\ \sigma(i - n) + n, & \text{otherwise.} \end{cases}$$

Example

If $\pi = 132$ and $\sigma = 4231$ then $\pi \oplus \sigma = 1327564$.

The direct sum is clearly associative.

Block number

Definition

A nonempty permutation which is not a direct sum of two nonempty permutations is called \oplus -irreducible.

Each permutation π can be written uniquely as a direct sum of \oplus -irreducible ones, called the **blocks** of π . Their number $\text{bl}(\pi)$ is the **block number** of π .

Example

$$\text{bl}(45321) = 1,$$

$$\text{bl}(312 \mid 54) = 2,$$

$$\text{bl}(1 \mid 2 \mid 3 \mid 4) = 4.$$

Remarks

- Direct sums and block decomposition of permutations appear naturally in the study of pattern-avoiding classes (Albert, Atkinson, Vatter).
- The block number of an arbitrary permutation was previously studied by Richard Stanley (2005), as the cardinality of the connectivity set (defined by Comtet).

Last descent

Definition

For a permutation $\pi \in \mathcal{S}_n$ let

$$\text{Ides}(\pi) := \max\{i : i \in \text{Des}(\pi)\},$$

with $\text{Ides}(\pi) := 0$ if $\text{Des}(\pi) = \emptyset$ (i.e., if π is the identity permutation).

Example

$$\text{Ides}(3176245) = 4$$

The sets $B_{l_{n,k}}$ and $L_{n,k}$

Definition

Let

$$B_{l_{n,k}} := \{\pi \in \mathcal{S}_n(321) : \text{bl}(\pi) = k\}.$$

Note that $\text{bl}(\pi) = \text{bl}(\pi^{-1})$.

Definition

Let

$$L_{n,k} = \{\pi \in \mathcal{S}_n(321) : \text{Ides}(\pi^{-1}) = k\}.$$

Cardinality

Definition

Recall: The n -th *Catalan number* is

$$C_n = \frac{1}{n+1} \binom{2n}{n}.$$

The corresponding generating function is

$$c(x) = \sum_{n=0}^{\infty} C_n x^n = \frac{1 - \sqrt{1 - 4x}}{2x}.$$

Cardinality

Definition

For each $k \geq 0$, the *n -th k -fold Catalan number* is the coefficient of x^n in $(xc(x))^k$. Explicitly:

$$C_{n,k} = \frac{k}{2n-k} \binom{2n-k}{n}.$$

Proposition

For positive integers $n \geq k \geq 1$:

$$C_{n,k} = |\text{SYT}(n-1, n-k)| = |L_{n,n-k}| = |B_{n,k}|$$

This result will be refined in the sequel.

Left-to-right maxima

Definition

The set of *left-to-right maxima* of $\pi \in \mathcal{S}_n$ is

$$\text{ltrMax}(\pi) = \{i \mid \pi(i) > \pi(j) \text{ for all } i < j\}$$

Example

$$\pi = \bar{3}12\bar{5}4\bar{6}.$$

Observation

For 321-avoiding permutations, the set of left-to-right maxima determines the descent set. Explicitly, for any $1 \leq i \leq n-1$,

$$i \in \text{Des}(\pi) \iff i \in \text{ltrMax}(\pi) \text{ and } i+1 \notin \text{ltrMax}(\pi).$$

Left-to-right maxima

Definition

The set of *left-to-right maxima* of $\pi \in \mathcal{S}_n$ is

$$\text{ltrMax}(\pi) = \{i \mid \pi(i) > \pi(j) \text{ for all } i < j\}$$

Example

$$\pi = \bar{3}12\bar{5}4\bar{6}.$$

Observation

For 321-avoiding permutations, the set of left-to-right maxima determines the descent set. Explicitly, for any $1 \leq i \leq n - 1$,

$$i \in \text{Des}(\pi) \iff i \in \text{ltrMax}(\pi) \text{ and } i + 1 \notin \text{ltrMax}(\pi).$$

Main result 1: Equi-distribution

Theorem (Adin-B.-Roichman '16)

For every positive integer n ,

$$\sum_{\pi \in \mathcal{S}_n(321)} \mathbf{x}^{\text{ltrMax}(\pi^{-1})} q^{\text{bl}(\pi)} = \sum_{\pi \in \mathcal{S}_n(321)} \mathbf{x}^{\text{ltrMax}(\pi^{-1})} q^{n - \text{lDes}(\pi)}.$$

Note the analogy with the classical

Theorem (Foata-Schützenberger '70)

For every positive integer n ,

$$\sum_{\pi \in \mathcal{S}_n} \mathbf{x}^{\text{Des}(\pi^{-1})} q^{\text{inv}(\pi)} = \sum_{\pi \in \mathcal{S}_n} \mathbf{x}^{\text{Des}(\pi^{-1})} q^{\text{maj}(\pi)}.$$

Main result 1: Equi-distribution

Theorem (Adin-B.-Roichman '16)

For every positive integer n ,

$$\sum_{\pi \in \mathcal{S}_n(321)} \mathbf{x}^{\text{ltrMax}(\pi^{-1})} q^{\text{bl}(\pi)} = \sum_{\pi \in \mathcal{S}_n(321)} \mathbf{x}^{\text{ltrMax}(\pi^{-1})} q^{n - \text{lDes}(\pi)}.$$

Note the analogy with the classical

Theorem (Foata-Schützenberger '70)

For every positive integer n ,

$$\sum_{\pi \in \mathcal{S}_n} \mathbf{x}^{\text{Des}(\pi^{-1})} q^{\text{inv}(\pi)} = \sum_{\pi \in \mathcal{S}_n} \mathbf{x}^{\text{Des}(\pi^{-1})} q^{\text{maj}(\pi)}.$$

Symmetry and Schur-positivity

Symmetric functions

Definition

A symmetric function is a formal power series $f \in \mathbb{C}[[x_1, x_2, \dots]]$ which is invariant under any permutation of the variables.

- We sometimes restrict to a finite number of variables by setting almost all of them to zero.

Example

$f = x_1 + x_2 + x_3$ is symmetric and homogeneous of degree 1. (with $x_4 = x_5 = \dots = 0$).

Semistandard tableaux

Definition

Let λ be a partition. A **semistandard Young tableau** of shape λ is a filling of the cells of λ by positive integers such that

- The entries in each row are **weakly increasing**.
- The entries in each column are **strictly increasing**.

Example

$$\lambda = (4, 3, 2) \quad T = \begin{array}{|c|c|c|c|} \hline 1 & 2 & 2 & 3 \\ \hline 3 & 3 & 4 & \\ \hline 5 & 6 & & \\ \hline \end{array}$$

Schur functions

With each semistandard Young tableau T we associate a monomial

$$\mathbf{x}^T = \prod_i x_i^{\text{number of } i\text{'s in } T}.$$

Example

$$T = \begin{array}{|c|c|c|c|} \hline 1 & 2 & 2 & 3 \\ \hline 3 & 3 & 4 & \\ \hline 5 & 6 & & \\ \hline \end{array}$$

$$\mathbf{x}^T = x_1 x_2^2 x_3^3 x_4 x_5 x_6.$$

The **Schur function** s_λ associated with a partition λ is defined by

$$s_\lambda = \sum_{T \in \text{SSYT}(\lambda)} \mathbf{x}^T.$$

Schur functions

Example

For $\lambda = (2, 1)$, the semistandard tableaux of shape λ filled with numbers out of $\{1, 2, 3\}$ are

$$\begin{array}{|c|c|} \hline 1 & 1 \\ \hline 2 & \\ \hline \end{array}, \begin{array}{|c|c|} \hline 1 & 1 \\ \hline 3 & \\ \hline \end{array}, \begin{array}{|c|c|} \hline 1 & 2 \\ \hline 2 & \\ \hline \end{array}, \begin{array}{|c|c|} \hline 1 & 2 \\ \hline 3 & \\ \hline \end{array}, \begin{array}{|c|c|} \hline 1 & 3 \\ \hline 2 & \\ \hline \end{array}, \begin{array}{|c|c|} \hline 1 & 3 \\ \hline 3 & \\ \hline \end{array}, \begin{array}{|c|c|} \hline 2 & 2 \\ \hline 3 & \\ \hline \end{array}, \begin{array}{|c|c|} \hline 2 & 3 \\ \hline 3 & \\ \hline \end{array}.$$

The corresponding Schur polynomial is

$$s_{(2,1)}(x_1, x_2, x_3) = x_1^2 x_2 + x_1^2 x_3 + x_1 x_2^2 + 2x_1 x_2 x_3 + x_1 x_3^2 + x_2^2 x_3 + x_2 x_3^2$$

Proposition

$\{s_\lambda \mid \lambda \vdash n\}$ is a basis for the vector space of symmetric functions which are homogeneous of degree n .

Schur functions

Example

For $\lambda = (2, 1)$, the semistandard tableaux of shape λ filled with numbers out of $\{1, 2, 3\}$ are

$$\begin{array}{|c|c|} \hline 1 & 1 \\ \hline 2 & \\ \hline \end{array}, \begin{array}{|c|c|} \hline 1 & 1 \\ \hline 3 & \\ \hline \end{array}, \begin{array}{|c|c|} \hline 1 & 2 \\ \hline 2 & \\ \hline \end{array}, \begin{array}{|c|c|} \hline 1 & 2 \\ \hline 3 & \\ \hline \end{array}, \begin{array}{|c|c|} \hline 1 & 3 \\ \hline 2 & \\ \hline \end{array}, \begin{array}{|c|c|} \hline 1 & 3 \\ \hline 3 & \\ \hline \end{array}, \begin{array}{|c|c|} \hline 2 & 2 \\ \hline 3 & \\ \hline \end{array}, \begin{array}{|c|c|} \hline 2 & 3 \\ \hline 3 & \\ \hline \end{array}.$$

The corresponding Schur polynomial is

$$s_{(2,1)}(x_1, x_2, x_3) = x_1^2 x_2 + x_1^2 x_3 + x_1 x_2^2 + 2x_1 x_2 x_3 + x_1 x_3^2 + x_2^2 x_3 + x_2 x_3^2$$

Proposition

$\{s_\lambda \mid \lambda \vdash n\}$ is a basis for the vector space of symmetric functions which are homogeneous of degree n .

Schur-positivity

Definition

A symmetric function is called **Schur-positive** if all the coefficients in its expansion in the basis of Schur functions are non-negative.

Example

For $\lambda \vdash k$ and $\mu \vdash \ell$, consider the product

$$s_\lambda s_\mu = \sum_{\nu} c_{\lambda, \mu}^{\nu} s_{\nu}.$$

The **Littlewood-Richardson rule** provides a combinatorial interpretation of the coefficients $c_{\lambda, \mu}^{\nu}$, proving that $s_\lambda s_\mu$ is Schur-positive.

An equivalent definition of symmetric functions

A formal power series $f(x_1, x_2, \dots)$ is **symmetric** if for every composition $\alpha = (\alpha_1, \dots, \alpha_n)$, all monomials $x_{i_1}^{\alpha_1} \cdots x_{i_k}^{\alpha_k}$ in f with distinct indices have the same coefficient.

Example

$$f = \sum_{i \neq j} x_i^3 x_j = x_1^3 x_2 + x_1 x_2^3 + x_1^3 x_3 + x_1 x_3^3 + x_2^3 x_3 + x_2 x_3^3 + \dots$$

Quasi-symmetric functions

A formal power series $f(x_1, x_2, \dots)$ is **quasi-symmetric** if for every composition $(\alpha_1, \dots, \alpha_k)$, all monomials $x_{i_1}^{\alpha_1} \cdots x_{i_k}^{\alpha_k}$ in f with indices $i_1 < i_2 < \dots < i_k$ have the same coefficients.

Example

$$f = \sum_{i < j} x_i^2 x_j = x_1^2 x_2 + x_1^2 x_3 + x_2^2 x_3 + \dots$$

is quasi-symmetric but not symmetric.

Denote by $QSym$ the vector space of quasi-symmetric functions which are homogeneous of degree n .

Quasi-symmetric functions

A formal power series $f(x_1, x_2, \dots)$ is **quasi-symmetric** if for every composition $(\alpha_1, \dots, \alpha_k)$, all monomials $x_{i_1}^{\alpha_1} \cdots x_{i_k}^{\alpha_k}$ in f with indices $i_1 < i_2 < \dots < i_k$ have the same coefficients.

Example

$$f = \sum_{i < j} x_i^2 x_j = x_1^2 x_2 + x_1^2 x_3 + x_2^2 x_3 + \dots$$

is quasi-symmetric but not symmetric.

Denote by $QSym$ the vector space of quasi-symmetric functions which are homogeneous of degree n .

The fundamental basis

For each subset $J \subseteq [n-1]$ define the corresponding **fundamental quasi-symmetric function**

$$F_J(\mathbf{x}) := \sum_{\substack{i_1 \leq i_2 \leq \dots \leq i_n \\ i_j < i_{j+1} \text{ if } j \in J}} x_{i_1} x_{i_2} \cdots x_{i_n}.$$

In particular, J can be the descent set of a permutation.

Example

$$\pi = 132, \text{Des}(\pi) = \{2\}.$$

$$\mathcal{F}_{\text{Des}(132)} = \mathcal{F}_{\{2\}} = x_1 x_1 x_2 + x_1 x_1 x_3 + x_1 x_2 x_3 + x_2 x_2 x_3 + \dots$$

Proposition (Gessel)

$\{F_J \mid J \subseteq [n-1]\}$ is a basis for $QSym_n$.

Schur-positivity

For $A \subseteq \mathcal{S}_n$, let

$$Q(A) = \sum_{\pi \in A} \mathcal{F}_{\text{Des}(\pi)}.$$

$Q(A)$ is called **Schur-positive** if it is symmetric and can be written as a linear combination of Schur functions with non-negative coefficients.

Question (Adin-Roichman, '13)

For which $A \subseteq \mathcal{S}_n$ is $Q(A)$ (symmetric and) Schur-positive?

Schur-positivity

For $A \subseteq \mathcal{S}_n$, let

$$Q(A) = \sum_{\pi \in A} \mathcal{F}_{\text{Des}(\pi)}.$$

$Q(A)$ is called **Schur-positive** if it is symmetric and can be written as a linear combination of Schur functions with non-negative coefficients.

Question (Adin-Roichman, '13)

For which $A \subseteq \mathcal{S}_n$ is $Q(A)$ (symmetric and) Schur-positive?

Main result 2: Schur-positivity of $Bl_{n,k}$

Recall

Definition

$$Bl_{n,k} := \{\pi \in \mathcal{S}_n(321) : \text{bl}(\pi) = k\}.$$

Theorem (Adin-B.-Roichman '16)

$\mathcal{Q}(Bl_{n,k})$ is (symmetric and) Schur positive.

Main result 2: Schur-positivity of $Bl_{n,k}$

Recall

Definition

$$Bl_{n,k} := \{\pi \in \mathcal{S}_n(321) : \text{bl}(\pi) = k\}.$$

Theorem (Adin-B.-Roichman '16)

$\mathcal{Q}(Bl_{n,k})$ is (symmetric and) Schur positive.

Main result 3: The character

Recall that the **Frobenius image** of an S_n -character $\chi = \sum_{\lambda \vdash n} c_\lambda \chi^\lambda$ is the symmetric function $f = \sum_{\lambda \vdash n} c_\lambda s_\lambda$, denoted by $ch(\chi)$.

Theorem (Adin-B.-Roichman '16)

For every positive integer $1 \leq k \leq n - 1$

$$Q(BI_{n,k}) = ch(\chi^{(n-1, n-k)} \downarrow_{S_n}^{S_{2n-k-1}})$$

and, for $k = n$,

$$Q(BI_{n,k}) = ch(\chi^{(n)}) = s_{(n)}.$$

Proof idea

Proof idea: bijection

The proofs use an explicit **left-to-right-maxima preserving** bijection from $Bl_{n,k}$ to $L_{n,n-k}$.

Definition

Define a map $f_n : \mathcal{S}_n(321) \mapsto \mathcal{S}_n(321)$, recursively on n , as follows. Each permutation $\pi \in \mathcal{S}_n$ belongs to exactly one of the following 3 classes, distinguished according to the location of the letter n and the relative order of $n-1$ and n .

- L : n is the **last** letter.
- D : n is not the last letter, and $n-1$ **precedes** n .
- R : $n-1$ is to the **right** of n .

Proof idea: bijection

The proofs use an explicit **left-to-right-maxima preserving** bijection from $Bl_{n,k}$ to $L_{n,n-k}$.

Definition

Define a map $f_n : \mathcal{S}_n(321) \mapsto \mathcal{S}_n(321)$, recursively on n , as follows. Each permutation $\pi \in \mathcal{S}_n$ belongs to exactly one of the following 3 classes, distinguished according to the location of the letter n and the relative order of $n-1$ and n .

- L : n is the **last** letter.
- D : n is not the last letter, and $n-1$ **precedes** n .
- R : $n-1$ is to the **right** of n .

Proof idea: bijection

Case L: n is the last letter.

- Omit n
- Apply f_{n-1} ;
- Insert n at the last position.

Case D: $n - 1$ is left of n , but n is not the last letter.

- Omit n .
- Apply f_{n-1} .
- Multiply from left by the transposition $(n - k - 1, n - k)$.
- Insert n at the same position as in π .

Case R: $n - 1$ is right of n .

In this case $n - 1$ must be the last letter.

- Exchange $n - 1$ and n in π , then omit n .
- Apply f_{n-1}
- Multiply (from the left) the resulting permutation by the cycle $(n - k, n - k + 1, \dots, n - 1, n)$.

Example

Let $\pi_8 = \pi = 31254786$.

$$\begin{array}{l}
 \pi_8 = 312 \mid 54 \mid 786 \xrightarrow[(45)]{D} \pi_7 = 3125476 \xrightarrow[(4567)]{R} \pi_6 = 312 \mid 54 \mid 6 \\
 \xrightarrow{L} \pi_5 = 312 \mid 54 \xrightarrow[(345)]{R} \pi_4 = 312 \mid 4 \\
 \xrightarrow{L} \pi_3 = 312 \xrightarrow[(23)]{R} \pi_2 = 21
 \end{array}$$

Example (cont.)

In the other direction:

$$\begin{aligned}
 f(\pi_2) = 21 & \xrightarrow{(23)} f(\pi_3) = 312 \rightarrow f(\pi_4) = 3124 \\
 & \xrightarrow{(345)} f(\pi_5) = 41253 \xrightarrow{(45)} f(\pi_6) = 412536 \\
 & \xrightarrow{(4567)} f(\pi_7) = 5126374 \xrightarrow{(45)} f(\pi_8) = 41263785
 \end{aligned}$$

Open problems

Open problems

- 1 Find a non-recursive definition for the bijection.
- 2 A **pattern-statistic pair** (Π, stat) consists of a subset $\Pi \subseteq \mathcal{S}_m$ and a permutation statistic $\text{stat} : \mathcal{S}_n \rightarrow \mathbb{N}$. It is **Schur-positive** if

$$Q(\{\pi \in \mathcal{S}_n(\Pi) \mid \text{stat}(\pi) = k\})$$

is Schur-positive for all positive integers n and k .

Find Schur-positive pattern-statistic pairs.

Open problems

- 1 Find a non-recursive definition for the bijection.
- 2 A **pattern-statistic pair** (Π, stat) consists of a subset $\Pi \subseteq \mathcal{S}_m$ and a permutation statistic $\text{stat} : \mathcal{S}_n \rightarrow \mathbb{N}$. It is **Schur-positive** if

$$\mathcal{Q}(\{\pi \in \mathcal{S}_n(\Pi) \mid \text{stat}(\pi) = k\})$$

is Schur-positive for all positive integers n and k .

Find Schur-positive pattern-statistic pairs.

Open problems

- 1 Find a non-recursive definition for the bijection.
- 2 A **pattern-statistic pair** (Π, stat) consists of a subset $\Pi \subseteq \mathcal{S}_m$ and a permutation statistic $\text{stat} : \mathcal{S}_n \rightarrow \mathbb{N}$. It is **Schur-positive** if

$$\mathcal{Q}(\{\pi \in \mathcal{S}_n(\Pi) \mid \text{stat}(\pi) = k\})$$

is Schur-positive for all positive integers n and k .

Find Schur-positive pattern-statistic pairs.

Thank you
for your attention!